

- [3] 1. Find the derivative directly from the definition for the function  $f(x) = \frac{1}{x+2}$ . You must use the definition, not some other method.
- [6] 2. Find the derivative of each function.
- a)  $f(x) = \sin(\ln(x^2))$
- b)  $g(t) = t^{\ln(t)}$
- [3] 3. Estimate the value of  $f(0.1)$  using a linearization of  $f(x) = \tan(x) + 1$ . Choose the point  $a$  of the linearization appropriately.
- [3] 4. Show that  $\frac{d}{dx} \left( \int_3^{x^2} \frac{1/2}{1+t} dt + \int_{\tan^{-1} x}^2 \tan(t) dt \right)$  is zero.
- [6] 5. Consider the curve defined by  $y + xe^y = x^2$ .
- a) Give  $\frac{dy}{dx}$  in terms of  $x$  and  $y$ .
- b) Find all values of  $x$  such that the point  $(x, 0)$  is on the curve defined by the above equation. For each of these give the slope of the tangent line to the curve at that point.
- [6] 6. Find each of the limits.
- a)  $\lim_{x \rightarrow 3^+} \frac{1 - e^{x-3}}{(x-3)^2}$
- b)  $\lim_{x \rightarrow 0} (x^2 + 1)^{1/x}$
- [4] 7. A ladder of length 2m is leaning against wall. The top of the ladder slides vertically down the wall while the bottom slides horizontally directly away from the wall.
- When the bottom of the ladder is 1m from the wall and moving at 0.1m/s, how fast is the top of the ladder falling?
- [6] 8. Evaluate each integral.
- a)  $\int xe^x dx$
- b)  $\int (\ln(x))^2 dx$

[6] 9. Evaluate each integral.

a)  $\int \frac{4}{(x+1)^2(x-1)} dx$

b)  $\int \frac{x}{\sqrt{(x+2)^2-1}} dx$

[8] 10. Evaluate each integral.

a)  $\int_0^1 \frac{e^x}{e^{2x} + 5e^x + 6} dx$

b)  $\int_0^{\pi/4} \frac{e^{\tan x}}{\cos^2 x} dx$

[4] 11. We wish to evaluate  $\int_2^3 \ln(x) dx$  numerically.

a) Give an expression for the Riemann sum for  $\int_2^3 \ln(x) dx$  using  $n = 3$  rectangles and the right-hand rule. You do not need to evaluate your expression numerically.

b) The difference between  $\int_a^b f(x) dx$  and the approximation using Simpson's method with  $n$  subintervals (or "rectangles") is at most  $\frac{K(b-a)^5}{180n^4}$ , where  $|f''''(x)| \leq K$  on  $[a, b]$ .

Give an expression for the value of  $n$  required so that Simpson's method applied to  $\int_2^3 \ln(x) dx$  is accurate to within 0.00001. You do not need to compute Simpson's method, nor evaluate your expression numerically. An expression for  $n$  suffices.

[5] 12. Consider the following function, and its derivatives.

$$f(x) = \frac{e^{-x}}{x^2}$$

$$f'(x) = \frac{-e^{-x}(x+2)}{x^3}$$

$$f''(x) = \frac{e^{-x}(x^2 + 4x + 6)}{x^4}$$

- Identify all horizontal and vertical asymptotes.
- Determine where it is increasing and where it is decreasing. Identify all extrema (local maximum and minimum).
- Determine where it is concave up and where it is concave down. Identify all inflection points.
- Sketch the function, labelling the extrema, inflection points, asymptotes and the intercepts.

- [+4] **13.** (bonus) Consider a rectangle of dimensions  $2x \times x$  and a square of dimensions  $y \times y$ .

If the sum of the perimeters of the rectangle and the square is  $\ell$ , find the value of  $x$  and  $y$  (in terms of  $\ell$ ) that minimize the sum of the areas of the rectangle and the square.