

MARK: PHY1321 PHY1331 Fall 2019 Assignment 1. Due Sept 20, 6:00PM

/24 LAST NAME (TYPED): \_\_\_\_\_ ASSIGNED NUMBER(BRIGHSPEACE)\_\_\_\_\_

TA:

**Pressure, Temperature, Ideal Gas Equation. Released: Sept 13 Due: Sept 20 6PM**


Each question is marked out of 4 points, In addition to this there are 4 points for presentation/aesthetics for the whole assignment.

- 1 HS teacher duplicated Torricelli's barometer using a mineral oil, of density  $1130 \text{ kg/m}^3$ , as the working liquid. What was the height  $h$  of the oil column for normal atmospheric pressure?

$$\text{Since } P_0 = \rho g h \quad h = \frac{P_0}{\rho g} = \frac{1.013 \times 10^5 \text{ Pa}}{(1130 \text{ kg/m}^3)(9.81 \text{ kg/ms}^2)} = 9.13 \text{ m}$$

- 2 A U-tube of uniform cross-sectional area, open to the atmosphere, is partially filled with mercury. Water is then poured into both arms. If the equilibrium configuration of the tube is as shown in, with  $h_2 = 0.80 \text{ cm}$ , determine the value of  $h_1$ .

Let  $h$  be the height of the water column added to the right side of the U-tube. Then when equilibrium is reached, the situation is as shown in the sketch at right. Now consider two points,  $A$  and  $B$  shown in the sketch, at the level of the water-mercury interface. By Pascal's Principle, the absolute pressure at  $B$  is the same as that at  $A$ .



But,

$$P_A = P_0 + \rho_w g h_1 + \rho_{Hg} g h_2 \text{ and}$$

$$P_B = P_0 + \rho_w g (h_1 + h + h_2).$$

Thus, from  $P_A = P_B$ ,  $\rho_w h_1 + \rho_w h + \rho_{Hg} h_2 = \rho_w h + \rho_{Hg} h_2$ , or

$$h_1 = \left( \frac{\rho_{Hg}}{\rho_{water}} - 1 \right) h_2 = (13.61)(0.8 \text{ cm}) = 10.08 \text{ cm}$$

- 3 a) Show that 1 mole of any gas at atmospheric pressure and at  $0^\circ \text{C}$  is taking 22.4 liters of volume  
 b) Show that the density of an ideal gas occupying a volume  $V$  is given by  $\rho = PM/RT$ , where  $M$  is the molar mass.  
 (c) Determine the density of oxygen gas at atmospheric pressure and  $20.0^\circ \text{C}$ .

a)  $pV = nRT \Rightarrow (101000)V = (1)(8.31)(273) = 0.0224 \text{ m}^3 = 22.4 \text{ l}$

b)  $pV = nRT \Rightarrow pV = \frac{m}{M} RT \Rightarrow \frac{pM}{RT} = \frac{m}{V} \Rightarrow \rho = \frac{pM}{RT}$

c)  $\rho = \frac{pM}{RT} = \frac{(101300)(0.032)}{(8.31)(293)} = 1.331 \frac{\text{kg}}{\text{m}^3}$

LAST NAME (TYPED): \_\_\_\_\_ ASSIGNED NUMBER(BRIGHSPEACE)\_\_\_\_\_

- 4      80 grams of oxygen and 40 grams hydrogen gas occupy separate, equal sections of 200 liter tank. The divide is removed and the gases are allowed to mix and react with each other. The temperature is kept constant at 110 °C, throughout the process.
- find the pressure of each gas in the separate containers .
  - find the pressure after the reaction ends.

There are 2.5moles of molecular oxygen gas in the first section, so

$$P_i V_i = nRT \Rightarrow P_i(O_2) = \frac{2.5 \cdot 8.31 \cdot 383}{0.1} Pa = Pa = 79568 Pa$$

There are 20 moles of molecular hydrogen gas in the first section, so

$$P_i V_i = nRT \Rightarrow P_i(H_2) = \frac{20 \cdot 8.31 \cdot 383}{0.1} Pa = 636546 Pa = 636 kPa$$

B) after the reaction there will be 5 moles of steam and 15moles of molecular hydrogen in a 200 liter container. The final pressure will be a sum of the partial pressures taken independently

$$P_f V_f = nRT \Rightarrow P_f(H_2O) = \frac{(5) (8.31) (383)}{0.2} Pa = 79568 Pa = 79.6 kPa$$

$$P_f V_f = nRT \Rightarrow P_f(H_2) = \frac{(15) (8.31) (383)}{0.2} Pa = 238705 = 239 kPa$$

- 5      A telescope forms an image of part of a cluster of stars on a square silicon charge-coupled detector (CCD) chip 2.00 cm on each side. A star field is focused on the CCD when it is first turned on and its temperature is 20.0°C. The star field contains 5 432 stars scattered uniformly. To make the detector more sensitive, it is cooled to -200°C. How many star images then fit onto the chip? The average coefficient of linear expansion of silicon is  $4.86 \times 10^{-6} (^\circ C)^{-1}$ .

$$\Delta S = \gamma S \Delta T = 2\alpha S \Delta T = 2(4.86 \cdot 10^{-6}) \cdot (4) \cdot (-220) cm^2 = -0.0085536 cm^2$$

$$S_f = S_i + \Delta S = 3.991446 cm^2$$

$$N_f = \frac{S_f}{S_i} N = \frac{3.991446}{4} 5432 = 5420.38 = 5420 \text{ full stars}$$