

**SOLUTION TO FINAL EXAM**

MAT 1322D, Fall 2012

Total = 53 marks

**1.** (4 marks) Determine whether each of the following improper integrals is convergent or divergent. If it is convergent, find its value; if it is divergent, use appropriate method to justify your conclusion:

(a)  $\int_0^{\infty} \frac{1}{(x+1)(x+3)} dx.$

*Solution.* Use partial fraction,  $\frac{1}{(x+1)(x+3)} = \frac{1}{2} \left( \frac{1}{x+1} - \frac{1}{x+3} \right).$

$$\begin{aligned} \int_0^{\infty} \frac{1}{(x+1)(x+3)} dx &= \frac{1}{2} \lim_{b \rightarrow \infty} \left( \int_0^b \frac{dx}{x+1} - \int_0^b \frac{dx}{x+3} \right) = \frac{1}{2} \lim_{b \rightarrow \infty} (\ln(b+1) - \ln(b+3) + \ln 3) \\ &= \frac{1}{2} \lim_{b \rightarrow \infty} \left( \ln \frac{b+1}{b+3} + \ln 3 \right) = \frac{1}{2} \ln 3. \end{aligned}$$

(b)  $\int_0^1 \frac{1}{\sqrt{x^2 + 2x^3}} dx.$

*Solution.* Since  $x^3 \leq x^2$  in the interval  $[0, 1]$ ,  $\frac{1}{\sqrt{x^2 + 2x^3}} > \frac{1}{\sqrt{3x^2}} = \frac{1}{\sqrt{3}x}.$  The improper integral

$$\int_0^1 \frac{1}{\sqrt{3}x} dx = \frac{1}{\sqrt{3}} \int_0^1 \frac{1}{x} dx \text{ is divergent, this integral is divergent.}$$

**2.** (4 marks) Find the volume of a solid whose base is the region on the  $x$ - $y$  plane bounded by the  $x$ -axis and the graph of the function  $y = 2(1 - x^2)$ . The cross sections of the solid perpendicular to the  $x$ -axis are semicircles with diameter on the  $x$ - $y$  plane. Find the volume of this solid.

*Solution.* The area of the cross section is  $A(x) = \frac{1}{2} \pi (1 - x^2)^2$ . The volume of the solid is


$$V = \frac{1}{2} \pi \int_{-1}^1 (1 - x^2)^2 dx = \pi \left[ x - \frac{2x^3}{3} + \frac{x^5}{5} \right]_{x=0}^1 = \frac{8}{15} \pi.$$

**3.** (4 marks) Find the length of the arc  $y = \frac{1}{2}x^2 - \frac{1}{4}\ln x$ ,  $1 \leq x \leq 2$ .

*Solution.*  $y' = x - \frac{1}{4x}$ ,  $(y')^2 = x^2 - \frac{1}{2} + \frac{1}{16x^2}$ .  $1 + (y')^2 = x^2 + \frac{1}{2} + \frac{1}{16x^2} = \left(x + \frac{1}{4x}\right)^2$ .

The length of the arc is  $L = \int_1^2 \left(x + \frac{1}{4x}\right) dx = \left[\frac{x^2}{2} + \frac{1}{4} \ln x\right]_{x=1}^2 = \frac{3}{2} + \frac{\ln 2}{4}$ .

**4.** (5 marks) A monument of the shape of a truncated circular cone is built with stone of density  $5000 \text{ kg/m}^3$ . The radius of the bottom of the monument is 3 meters, the radius of the top is 1 meter, and the height of the monument is 20 meters. (Recall that the radius of the cross section at height  $h$  is  $3 - h/10$  meters). Find the work (in Joules) needed to lift the stone from the ground level to build the monument. (Use  $g = 9.8 \text{ m/sec}^2$ )



*Solution.* A layer of the monument at height  $h$  of thickness  $dh$  weighs

$$dw = \delta g \pi (3 - h/10)^2 dh.$$


The work needed to lift the stone of this layer from the ground to height  $h$  is

$$dW = \delta g \pi (3 - h/10)^2 h dh.$$

The total work needed is

$$W = \delta g \pi \int_0^{20} \left(3 - \frac{h}{10}\right)^2 h dh = \delta g \pi \left[ \frac{9}{2} h^2 - \frac{1}{5} h^3 + \frac{1}{400} h^4 \right]_{h=0}^{20} \approx 2.94 \times 10^7 \text{ Joule.}$$

**5.** (4 marks) Suppose that a triangular surface is submerged into water as shown in the following figure. The top of the triangle is 2 meters under the water surface. The height of the triangle is 3 meters, and the length of the top side is 4 meters. Find the force acting on this surface.



*Solution.* Take a slice of the surface at depth  $2 + D$  of thickness  $dD$ . The length of this slice is approximately  $x = \frac{4(3-D)}{3}$ . The force acting on this slice is

$$dF = 1000g \frac{4(3-D)(2+D)}{3} dD.$$

The total force is

$$F = \frac{4000g}{3} \int_0^3 (3-D)(2+D) dD = 18000g \approx 176400 \text{ Newton.}$$

**6.** (5 marks) Consider the initial-value problem:  $y' = 2t \cos^2 y$ ,  $y(1) = \pi/4$ .

(i) (3 marks) Solve this initial-value problem.

(ii) (2 marks) Use Euler's method with step size  $h = 0.25$  to find an approximation of  $y(1.5)$ . (Use 4 digits after the decimal point in your calculation).

*Solution.* (i) Separating variables,  $\int \frac{dy}{\cos^2 y} = \int 2t dt$ .  $\tan y = t^2 + C$ . By the initial condition  $C = 0$ . Then  $y = \arctan(t^2)$ .

(ii)  $t_0 = 1$ ,  $y(1) = y_0 = \pi/4 \approx 0.7854$ .

$t_1 = 0.25$ ,  $y(1.25) \approx y_1 = y_0 + h(2t_0 \cos^2 y_0) \approx 1.0354$ .

$t_2 = 0.50$ ,  $y(1.50) \approx y_2 = y_1 + h(2t_1 \cos^2 y_1) \approx 1.1981$ .

**7.** (6 marks) Consider the equation  $\frac{dy}{dt} = y(5-y)$ ,  $y(0) = 2$ .

(i) (3 marks) Solve this equation analytically.

(ii) (1 mark) Find the limit of the solution when  $t$  approaches infinity.

(iii) (2 marks) Sketch the graph of the solution of this initial-value problem. Mark the inflection point of the graph.


*Solution.* (i) Separating variables,  $\int \frac{dy}{y(5-y)} = \int dt$ . Use partial fraction,

$$\int \frac{dy}{y(5-y)} = \frac{1}{5} \left( \int \frac{1}{y} dy + \int \frac{1}{5-y} dy \right) = \frac{1}{5} \ln \left| \frac{y}{5-y} \right| = t + C.$$

$$\frac{y}{5-y} = Ke^{5t}. \text{ By the initial condition, } K = \frac{2}{3}. \text{ Hence, } y = \frac{10e^{5t}}{3+2e^{5t}} = \frac{10}{2+3e^{-5t}}.$$

(ii) The limit of the solution is  $10 / 2 = 5$ .

(iii) The graph of the solution looks like the following



**8. (6 marks)** Determine whether each of the following series is convergent or divergent. Justify your answer by appropriate test method and state the condition to use these tests:

(a)  $\sum_{n=1}^{\infty} \frac{n}{e^n}$ ;      (b)  $\sum_{n=2}^{\infty} \frac{(-1)^{n-1}}{\ln n}$ ;      (c)  $\sum_{n=1}^{\infty} \frac{2n-1}{\sqrt{n^3+n}}$ .

*Solution.* (a) Since the general term is positive, decreasing and continuous, we can use the integral test.

$$\int_0^{\infty} xe^{-x} dx = \lim_{b \rightarrow \infty} \left[ -(1+x)e^{-x} \right]_{x=0}^b = \lim_{b \rightarrow \infty} (1 - (b+1)e^{-b}) = 1 < \infty, \text{ this series is convergent.}$$

Since the series is positive, we may also use the ratio test:

$$\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \frac{(n+1)e^n}{ne^{n+1}} = \lim_{n \rightarrow \infty} \frac{1}{e} < 1. \text{ By the ratio test, this series converges.}$$

(b) Since the general term is alternating and decreasing, we can use the alternating series test. Since the general term approaches 0, this series is convergent.

(c) Since this is a positive series, we can use the limit comparison test.

Compare this series with the series  $\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}}$ . We have

$$\lim_{n \rightarrow \infty} \frac{2n-1}{\sqrt{n^3+n}} \sqrt{n} = \lim_{n \rightarrow \infty} \frac{\sqrt{n}(2n-1)/(n\sqrt{n})}{\sqrt{n^3+n}/(n\sqrt{n})} = \lim_{n \rightarrow \infty} \frac{2-1/n}{\sqrt{1+1/n^2}} = 2.$$

Since  $\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}}$  is a  $p$ -series with  $p = 1/2$ , it is divergent. Series  $\sum_{n=1}^{\infty} \frac{2n-1}{\sqrt{n^3+n}}$  is also divergent.

9. (4 marks) Find the radius of convergence and the interval of convergence of the power series

$$\sum_{n=1}^{\infty} \frac{(x+5)^n}{3^n n}.$$

*Solution.* The center of the series is  $-5$ . The radius of convergence is

$$\lim_{n \rightarrow \infty} \left| \frac{c_n}{c_{n+1}} \right| = \lim_{n \rightarrow \infty} \left| \frac{3^{n+1}(n+1)}{3^n n} \right| = 3. \quad \text{This series is absolutely convergent in the interval } (-8, -2).$$

When  $x = -8$ , the series is  $\sum_{n=1}^{\infty} \frac{(-3)^n}{3^n n} = \sum_{n=1}^{\infty} \frac{(-1)^n}{n}$ . This is an alternating decreasing series with general term approaching 0. By the alternating series test, this series is convergent.

When  $x = -2$ , the series is  $\sum_{n=1}^{\infty} \frac{3^n}{3^n n} = \sum_{n=1}^{\infty} \frac{1}{n}$ . It is the harmonic series, which is divergent. The interval of convergence of this series is  $[-8, -2)$ .

10. (5 marks) Recall that the binomial series is

$$(1+x)^k = \sum_{n=0}^{\infty} \frac{k(k-1)(k-2)\dots(k-n+1)}{n!} x^n = 1 + kx + \frac{k(k-1)}{2!} x^2 + \frac{k(k-1)(k-2)}{3!} x^3 + \dots$$

(i) (2 marks) Find the Maclaurin series of the function  $y = \frac{1}{\sqrt{1+x^2}}$ .

(ii) (3 marks) Use the fact that  $\frac{d}{dx} \ln(x + \sqrt{1+x^2}) = \frac{1}{x + \sqrt{1+x^2}} \left( 1 + \frac{x}{\sqrt{1+x^2}} \right) = \frac{1}{\sqrt{1+x^2}}$  to find the Maclaurin series of the function  $y = \ln(x + \sqrt{1+x^2})$ .

You may use or not use the sigma notation. If you don't use the sigma notation, give enough number of terms to demonstrate the answer.

$$\begin{aligned}
 \text{Solution. } \frac{1}{\sqrt{1+x^2}} &= (1+x^2)^{-1/2} = \sum_{n=0}^{\infty} \frac{\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)\cdots\left(-\frac{2n-1}{2}\right)}{n!} x^{2n} \\
 &= 1 + \left(-\frac{1}{2}\right)x^2 + \frac{1}{2!}\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)x^4 + \frac{1}{3!}\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)\left(-\frac{5}{2}\right)x^6 + \dots \\
 &= 1 - \frac{1}{2}x^2 + \frac{1 \cdot 3}{2!2^2}x^4 - \frac{1 \cdot 3 \cdot 5}{3!2^3}x^6 + \dots \\
 &= \sum_{n=0}^{\infty} \frac{1 \cdot 3 \cdot \dots \cdot (2n-1)}{n!2^n} x^{2n}.
 \end{aligned}$$

$$\begin{aligned}
 \ln(x + \sqrt{1+x^2}) &= \sum_{n=0}^{\infty} \frac{1 \cdot 3 \cdot \dots \cdot (2n-1)}{n!2^n(2n+1)} x^{2n+1} \\
 &= x - \frac{1}{2 \cdot 3}x^3 + \frac{1 \cdot 3}{2!2^2 \cdot 5}x^5 - \frac{1 \cdot 3 \cdot 5}{3!2^3 \cdot 7}x^7 + \dots = x - \frac{1}{6}x^3 + \frac{3}{40}x^5 - \frac{5}{112}x^7 + \dots
 \end{aligned}$$

**11.** (6 marks) Consider the 2-variable function  $z = f(x, y) = \frac{xy}{2x^2 + y^2 + 1}$ .

(a) (2 mark) Find the gradient vector of this function at the point (2, 1).

(b) (2 marks) Find the directional derivative of  $z$  at point (2, 1) in the direction of the vector  $\mathbf{v} = 3\mathbf{i} + 4\mathbf{j}$ .

(c) (2 marks) Find the equation of the plane tangent to the graph of this function at a point where  $x = 2$ , and  $y = 1$ .

$$\text{Solution. (a) } z_x = \frac{y(2x^2 + y^2 + 1) - 4x^2y}{(2x^2 + y^2 + 1)^2} = \frac{y^3 - 2x^2y + y}{(2x^2 + y^2 + 1)^2}.$$

$$z_y = \frac{x(2x^2 + y^2 + 1) - 2xy^2}{(2x^2 + y^2 + 1)^2} = \frac{2x^3 - xy^2 + x}{(2x^2 + y^2 + 1)^2}.$$

$$\text{When } x = 2, y = 1, z_x = \frac{-6}{100} = -\frac{3}{50}, \text{ and } z_y = \frac{16}{100} = \frac{4}{25}.$$

(c) The unit vector in the direction  $\mathbf{v}$  is  $\mathbf{u} = \frac{3}{5}\mathbf{i} + \frac{4}{5}\mathbf{j}$ . The directional derivative is

$$D_{\mathbf{v}}(z) = -\frac{6}{100} \times \frac{3}{5} + \frac{16}{100} \times \frac{4}{5} = \frac{46}{500} = \frac{23}{250}.$$

(d) When  $x = 2$ , and  $y = 1$ ,  $z = \frac{1}{5}$ . The equation of the tangent plane is

$$z = -\frac{3}{50}(x-2) + \frac{4}{25}(y-1) + \frac{1}{5}, \text{ or } 50z = -3x + 8y + 8, 3x - 8y + 50z = 8.$$